

Yersinia isolates confirmed by ERL in 2018

<i>Yersinia</i>	DNII																			Total		
	Auckland	Bay of Plenty	Canterbury	Capital and Coast	Combined Auckland	Combined Wellington	Courties Manukau	Hawkes Bay	Hutt Valley	Lakes	MidCentral	Nelson Marlborough	Northland	South Canterbury	Southern	Tairāwhiti	Taranaki	Waikato	Wairarapa		Waitemata	West Coast
<i>enterocolitica</i> *			2							1			1									4
<i>enterocolitica</i> biotype 1A	16	9	50	2	21	18	8	4		3		9		2	31		3	15	1	4	1	201
<i>enterocolitica</i> biotype 2/O serotype O:3					1																	1
<i>enterocolitica</i> biotype 2/O serotype O:5, 27	2	1	12	1	11	3				1		1			6		1	2				42
<i>enterocolitica</i> biotype 2/O serotype O:9	15	29	109	9	173	71	4	15	1	22	10	21	3	3	45	3	15	36	2	6	3	595
<i>enterocolitica</i> biotype 4 serotype O:3	3	25	13	2	22	40	2	5		15	5				5		3	17	2	3		207
<i>pseudotuberculosis</i>	1		7			3								1	2		1					15
Total	37	64	193	14	278	135	14	24	1	41	18	31	6	6	88	3	20	70	5	13	4	1063

*Species identification confirmed by whole genome sequencing analysis.